


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

|  |  |  |  |
|--|--|--|--|
| (51) International Patent Classification <sup>6</sup> :<br><b>A61K 7/48, 31/40, 31/44</b>  |  | <b>A1</b>  | (11) International Publication Number: <b>WO 98/55091</b>        |
|  |  |  | (43) International Publication Date: 10 December 1998 (10.12.98) |
| (1) International Application Number: <b>PCT/US98/11246</b><br>(22) International Filing Date: <b>3 June 1998 (03.06.98)</b><br>(30) Priority Data:<br><b>08/869,426</b> <b>4 June 1997 (04.06.97)</b> <b>US</b><br>(71) Applicant: <b>GUILFORD PHARMACEUTICALS INC.</b><br><b>[US/US]; 6611 Tributary Street, Baltimore, MD 21224 (US).</b><br>(72) Inventors: <b>HAMILTON, Gregory, S.; 6501 Frederick Road, Catonsville, MD 21228 (US). STEINER, Joseph, P.; 988 Sugar Maple Street, Hampstead, MD 21074 (US).</b><br>(74) Agent: <b>NATH, Gary, M.; Nath &amp; Associates, 6th floor, 1030 15th Street, N.W., Washington, DC 20005 (US).</b> |  | (81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b><br><br><b>Published</b><br><i>With international search report.</i><br><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |  |

(54) Title: **PYRROLIDINE DERIVATIVE HAIR GROWTH COMPOSITIONS AND USES**



(57) Abstract

This invention relates to pharmaceutical compositions and methods for treating alopecia and promoting hair growth using pyrrolidine derivatives.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

|    |                          |    |  |    |  |    |                          |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania                  | ES | Spain                                    | LS | Lesotho                                      | SI | Slovenia                 |
| AM | Armenia                  | FI | Finland                                  | LT | Lithuania                                    | SK | Slovakia                 |
| AT | Austria                  | FR | France                                   | LU | Luxembourg                                   | SN | Senegal                  |
| AU | Australia                | GA | Gabon                                    | LV | Latvia                                       | SZ | Swaziland                |
| AZ | Azerbaijan               | GB | United Kingdom                           | MC | Monaco                                       | TD | Chad                     |
| BA | Bosnia and Herzegovina   | GE | Georgia                                  | MD | Republic of Moldova                          | TG | Togo                     |
| BB | Barbados                 | GH | Ghana                                    | MG | Madagascar                                   | TJ | Tajikistan               |
| BE | Belgium                  | GN | Guinea                                   | MK | The former Yugoslav<br>Republic of Macedonia | TM | Turkmenistan             |
| BF | Burkina Faso             | GR | Greece                                   | ML | Mali   | TR | Turkey                   |
| BG | Bulgaria                 | HU | Hungary                                  | MN | Mongolia                                     | TT | Trinidad and Tobago      |
| BJ | Benin                    | IE | Ireland                                  | MR | Mauritania                                   | UA | Ukraine                  |
| BR | Brazil                   | IL | Israel                                   | MW | Malawi                                       | UG | Uganda                   |
| BY | Belarus                  | IS | Iceland                                  | MX | Mexico                                       | US | United States of America |
| CA | Canada                   | IT | Italy                                    | NE | Niger  | UZ | Uzbekistan               |
| CF | Central African Republic | JP | Japan                                    | NL | Netherlands                                  | VN | Viet Nam                 |
| CG | Congo                    | KE | Kenya                                    | NO | Norway                                       | YU | Yugoslavia               |
| CH | Switzerland              | KG | Kyrgyzstan                               | NZ | New Zealand                                  | ZW | Zimbabwe                 |
| CI | Côte d'Ivoire            | KP | Democratic People's<br>Republic of Korea | PL | Poland                                       |    |                          |
| CM | Cameroon                 | KR | Republic of Korea                        | PT | Portugal                                     |    |                          |
| CN | China                    | KZ | Kazakhstan                               | RO | Romania                                      |    |                          |
| CU | Cuba                     | LC | Saint Lucia                              | RU | Russian Federation                           |    |                          |
| CZ | Czech Republic           | LI | Liechtenstein                            | SD | Sudan  |    |                          |
| DE | Germany                  | LK | Sri Lanka                                | SE | Sweden                                       |    |                          |
| DK | Denmark                  | LR | Liberia                                  | SG | Singapore                                    |    |                          |
| EE | Estonia                  |    |  |    |  |    |                          |

## PYRROLIDINE DERIVATIVE

### HAIR GROWTH COMPOSITIONS AND USES

5 This application is a continuation-in-part of  
U.S. Patent Application No. 08/869,426, filed on June  
4, 1997, the entire contents of which are herein  
incorporated by reference.

### BACKGROUND OF THE INVENTION

10

#### 1. Field of Invention

This invention relates to pharmaceutical  
compositions and methods for treating alopecia and  
promoting hair growth using low molecular weight,  
15 small molecule pyrrolidine derivatives.

#### 2. Description of Related Art

Hair loss occurs in a variety of situations.  
These situations include male pattern alopecia,  
20 alopecia senilis, alopecia areata, diseases  
accompanied by basic skin lesions or tumors, and  
systematic disorders such as nutritional disorders and  
internal secretion disorders. The mechanisms causing  
hair loss are very complicated, but in some instances  
25 can be attributed to aging, genetic disposition, the  
activation of male hormones, the loss of blood supply  
to hair follicles, and scalp abnormalities.

The immunosuppressant drugs FK506, rapamycin and  
cyclosporin are well known as potent T-cell specific

immunosuppressants, and are effective against graft rejection after organ transplantation. It has been reported that topical, but not oral, application of FK506 (Yamamoto et al., J. Invest. Dermatol., 1994, 102, 160-164; Jiang et al., J. Invest. Dermatol. 1995, 104, 523-525) and cyclosporin (Iwabuchi et al., J. Dermatol. Sci. 1995, 9, 64-69) stimulates hair growth in a dose-dependent manner. One form of hair loss, alopecia areata, is known to be associated with autoimmune activities; hence, topically administered immunomodulatory compounds are expected to demonstrate efficacy for treating that type of hair loss. The hair growth stimulating effects of FK506 have been the subject of an international patent filing covering FK506 and structures related thereto for hair growth stimulation (Honbo et al., EP 0 423 714 A2). Honbo et al. discloses the use of relatively large tricyclic compounds, known for their immunosuppressive effects, as hair revitalizing agents.

The hair growth and revitalization effects of FK506 and related agents are disclosed in many U.S. patents (Goulet et al., U.S. Patent No. 5,258,389; Luly et al., U.S. Patent No. 5,457,111; Goulet et al., U.S. Patent No. 5,532,248; Goulet et al., U.S. Patent No. 5,189,042; and Ok et al., U.S. Patent No. 5,208,241; Rupprecht et al., U.S. Patent No. 5,284,840; Organ et al., U.S. Patent No. 5,284,877). These patents claim FK506 related compounds. Although

they do not claim methods of hair revitalization, they disclose the known use of FK506 for effecting hair growth. Similar to FK506 (and the claimed variations in the Honbo et al. patent), the compounds claimed in these patents are relatively large. Further, the cited patents relate to immunomodulatory compounds for use in autoimmune related diseases, for which FK506's efficacy is well known.

Other U.S. patents disclose the use of cyclosporin and related compounds for hair revitalization (Hauer et al., U.S. Patent No. 5,342,625; Eberle, U.S. Patent No. 5,284,826; Hewitt et al., U.S. Patent No. 4,996,193). These patents also relate to compounds useful for treating autoimmune diseases and cite the known use of cyclosporin and related immunosuppressive compounds for hair growth.

However, immunosuppressive compounds by definition suppress the immune system and also exhibit other toxic side effects. Accordingly, there is a need for non-immunosuppressant, small molecule compounds which are useful as hair revitalizing compounds.

Hamilton and Steiner disclose in U.S. Patent No. 5,614,547 novel pyrrolidine carboxylate compounds which bind to the immunophilin FKBP12 and stimulate nerve growth, but which lack immunosuppressive effects. Unexpectedly, it has been discovered that

these non-immunosuppressant compounds promote hair growth with an efficacy similar to FK506. Yet their novel small molecule structure and non-immunosuppressive properties differentiate them from FK506 and related immunosuppressive compounds found in the prior art.

#### SUMMARY OF THE INVENTION

The present invention relates to a method for treating alopecia or promoting hair growth in an animal, which comprises administering to said animal an effective amount of a pyrrolidine derivative.

The present invention further relates to a pharmaceutical composition which comprises:

(i) an effective amount of a pyrrolidine derivative for treating alopecia or promoting hair growth in an animal; and

(ii) a pharmaceutically acceptable carrier.

The pyrrolidine derivatives used in the inventive methods and pharmaceutical compositions have an affinity for FKBP-type immunophilins and do not exert any significant immunosuppressive activity.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a photograph of C57 Black 6 mice before being shaved for the hair regeneration experiment.

FIG. 2 is a photograph of mice treated with a vehicle after six weeks. FIG. 2 shows that less than

3% of the shaved area is covered with new hair growth when the vehicle (control) is administered.

FIG. 3 is a photograph of mice treated with 10  $\mu$ M of GPI 1046, one of the non-immunosuppressive pyrrolidine derivative neuroimmunophilin FKBP ligands of this application, after six weeks. FIG. 3 shows the remarkable effects of non-immunosuppressive neuro-immunophilin FKBP ligands, wherein 90% of the shaved area is covered with new hair growth.

FIG. 4 is a photograph of mice treated with 30  $\mu$ M of GPI 1046 after six weeks. FIG. 4 shows the remarkable ability of non-immunosuppressive neuroimmunophilin FKBP ligands to achieve, essentially, complete hair regrowth in the shaved area.

FIG. 5 is a bar graph depicting the relative hair growth indices for C57 Black 6 mice treated with a vehicle, FK506, and various non-immunosuppressive neuroimmunophilin FKBP ligands, including GPI 1046, 14 days after treatment with each identified compound. Figure 5 demonstrates the remarkable early hair growth promoted by a wide variety of non-immunosuppressive neuroimmunophilin FKBP ligands.

## DETAILED DESCRIPTION OF THE INVENTION

### Definitions


"Alopecia" refers to deficient hair growth and partial or complete loss of hair, including without

limitation androgenic alopecia (male pattern baldness), toxic alopecia, alopecia senilis, alopecia areata, alopecia pelada and trichotillomania. Alopecia results when the pilar cycle is disturbed.

5 The most frequent phenomenon is a shortening of the hair growth or anagen phase due to cessation of cell proliferation. This results in an early onset of the catagen phase, and consequently a large number of hairs in the telogen phase during which the follicles  
10 are detached from the dermal papillae, and the hairs fall out. Alopecia has a number of etiologies, including genetic factors, aging, local and systemic diseases, febrile conditions, mental stresses, hormonal problems, and secondary effects of drugs.

15 "GPI 1605" refers to a compound of formula


20



GPI 1605

"GPI 1046" refers to 3-(3-pyridyl)-1-propyl (2s)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, a compound of formula

25




GPI1046



"GPI 1312" refers to a compound of formula


5



GPI 1312

"GPI 1572" refers to a compound of formula


10



GPI 1572

"GPI 1389" refers to a compound of formula


15



GPI 1389

"GPI 1511" refers to a compound of formula


20



GPI 1511

25

"GPI 1234" refers to a compound of formula



GPI 1234

"Isomers" refer to different compounds that have the same molecular formula. "Stereoisomers" are isomers that differ only in the way the atoms are arranged in space. "Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other. "Diastereoisomers" are stereoisomers which are not mirror images of each other. "Racemic mixture" means a mixture containing equal parts of individual enantiomers. "Non-racemic mixture" is a mixture containing unequal parts of individual enantiomers or stereoisomers.

"Pharmaceutically acceptable salt, ester, or solvate" refers to a salt, ester, or solvate of a subject compound which possesses the desired pharmacological activity and which is neither biologically nor otherwise undesirable. A salt, ester, or solvate can be formed with inorganic acids such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, gluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, naphthylate, 2-naphthalenesulfonate, nicotinate, oxalate, sulfate, thiocyanate, tosylate and undecanoate. Examples of base salts, esters, or

solvates include ammonium salts; alkali metal salts, such as sodium and potassium salts; alkaline earth metal salts, such as calcium and magnesium salts; salts with organic bases, such as dicyclohexylamine salts; N-methyl-D-glucamine; and salts with amino acids, such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups can be quarternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides; dialkyl sulfates, such as dimethyl, diethyl, dibutyl, and diamyl sulfates; long chain halides, such as decyl, lauryl, myristyl, and stearyl chlorides, bromides, and iodides; aralkyl halides, such as benzyl and phenethyl bromides; and others. Water or oil-soluble or dispersible products are thereby obtained.

"Pilar cycle" refers to the life cycle of hair follicles, and includes three phases:

- (1) the anagen phase, the period of active hair growth which, insofar as scalp hair is concerned, lasts about three to five years;
- (2) the catagen phase, the period when growth stops and the follicle atrophies which, insofar as scalp hair is concerned, lasts about one to two weeks; and
- (3) the telogen phase, the rest period when hair progressively separates and finally falls out which, insofar as scalp hair is

concerned, lasts about three to four months. Normally 80 to 90 percent of the follicles are in the anagen phase, less than 1 percent being in the catagen phase, and the rest being in the telogen phase. In the telogen phase, hair is uniform in diameter with a slightly bulbous, non-pigmented root. By contrast, in the anagen phase, hair has a large colored bulb at its root.

"Promoting hair growth" refers to maintaining, inducing, stimulating, accelerating, or revitalizing the germination of hair.

"Treating alopecia" refers to:

(i) preventing alopecia in an animal which may be predisposed to alopecia; and/or

(ii) inhibiting, retarding or reducing alopecia; and/or

(iii) promoting hair growth; and/or

(iv) prolonging the anagen phase of the hair cycle; and/or

(v) converting vellus hair to growth as terminal hair. Terminal hair is coarse, pigmented, long hair in which the bulb of the hair follicle is seated deep in the dermis. Vellus hair, on the other hand, is fine, thin, non-pigmented short hair in which the hair bulb is located superficially in the dermis. As alopecia progresses, the hairs change from the terminal to the vellus type.

### Methods of the Present Invention

The present invention relates to a method for treating alopecia or promoting hair growth in an animal, which comprises administering to said animal  
5 an effective amount of a pyrrolidine derivative.

The inventive method is particularly useful for treating male pattern alopecia, alopecia senilis, alopecia areata, alopecia resulting from skin lesions or tumors, alopecia resulting from cancer therapy such  
10 as chemotherapy and radiation, and alopecia resulting from systematic disorders such as nutritional disorders and internal secretion disorders.

### Pharmaceutical Compositions of the Present Invention

15 The present invention also relates to a pharmaceutical composition comprising:

- (i) an effective amount of a pyrrolidine derivative for treating alopecia or promoting hair growth in an animal; and
- 20 (ii) a pharmaceutically acceptable carrier.

### PYRROLIDINE DERIVATIVES


The pyrrolidine derivatives used in the methods and pharmaceutical compositions of the present  
25 invention are low molecular weight, small molecule compounds having an affinity for an FKBP-type immunophilins, such as FKBP12. When a pyrrolidine derivative binds to an FKBP-type immunophilin, it has

been found to inhibit the prolyl peptidyl cis trans isomerase, or rotamase, activity of the binding protein. Unexpectedly, these compounds have also been found to stimulate hair growth. The compounds are  
5 devoid of any significant immunosuppressive activity.

FORMULA I

The pyrrolidine derivative may be a compound of formula I

10



15

or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

R<sub>1</sub> is C<sub>1</sub>-C<sub>8</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>8</sub> straight or branched chain alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl or Ar<sub>1</sub>, wherein said R<sub>1</sub> is  
20 unsubstituted or substituted with one or more substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, hydroxy, and Ar<sub>2</sub>;

25


Ar<sub>1</sub> and Ar<sub>2</sub> are independently selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl and phenyl, wherein said

Ar<sub>1</sub> is unsubstituted or substituted with one or more substituent(s) independently selected from the group consisting of hydrogen, halo, hydroxy, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, S, CH<sub>2</sub> or H<sub>2</sub>;

Y is O or NR<sub>2</sub>, wherein R<sub>2</sub> is hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; and

Z is C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, wherein said Z is substituted with one or more substituent(s) independently selected from the group consisting of Ar<sub>1</sub>, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl; or Z is fragment



5

wherein:

R<sub>3</sub> is C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl which is unsubstituted or substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl or Ar<sub>1</sub>;

10 X<sub>2</sub> is O or NR<sub>5</sub>, wherein R<sub>5</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, and C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl; and

15 R<sub>4</sub> is selected from the group consisting of phenyl, benzyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl substituted with phenyl, and C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl substituted with phenyl.

20 In a preferred embodiment of formula I, Z and R<sub>1</sub> are lipophilic.

In a more preferred embodiment of formula I, the compound is selected from the group consisting of:

25 3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-phenyl-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(3,4,5-trimethoxyphenyl)-1-propyl (2S)-1-(3,3-



dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(3,4,5-trimethoxyphenyl)-1-prop-2-(E)-enyl  
(2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-  
pyrrolidinecarboxylate;

5 3-(4,5-dichlorophenyl)-1-propyl (2S)-1-(3,3-  
dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(4,5-dichlorophenyl)-1-prop-2-(E)-enyl (2S)-1-  
(3,3-dimethyl-1,2-dioxopentyl)-2-  
pyrrolidinecarboxylate;

10 3-(4,5-methylenedioxyphenyl)-1-propyl (2S)-1-  
(3,3-dimethyl-1,2-dioxopentyl)-2-  
pyrrolidinecarboxylate;

3-(4,5-methylenedioxyphenyl)-1-prop-2-(E)-enyl  
(2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-  
15 pyrrolidinecarboxylate;

3-cyclohexyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-  
dioxopentyl)-2-pyrrolidinecarboxylate;

3-cyclohexyl-1-prop-2-(E)-enyl (2S)-1-(3,3-  
dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

20 (1R)-1,3-diphenyl-1-propyl (2S)-1-(3,3-dimethyl-  
1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

(1R)-1,3-diphenyl-1-prop-2-(E)-enyl (2S)-1-(3,3-  
dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

(1R)-1-cyclohexyl-3-phenyl-1-propyl (2S)-1-(3,3-  
25 dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

(1R)-1-cyclohexyl-3-phenyl-1-prop-2-(E)-enyl  
(2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-  
pyrrolidinecarboxylate;

- 5      cyclonhexyl)ethyl-2-pyrrolidinecarboxylate;
- 3-phenyl-1-propyl      (2*S*)-1-(1,2-dioxo-4-cyclohexyl)butyl-2-pyrrolidinecarboxylate;
- 3-phenyl-1-propyl      (2*S*)-1-(1,2-dioxo-2-[2-furanyl])ethyl-2-pyrrolidinecarboxylate;
- 10      3-phenyl-1-propyl      (2*S*)-1-(1,2-dioxo-2-[2-thienyl])ethyl-2-pyrrolidinecarboxylate;
- 3-phenyl-1-propyl      (2*S*)-1-(1,2-dioxo-2-[2-thiazolyl])ethyl-2-pyrrolidinecarboxylate;
- 15      3-phenyl-1-propyl      (2*S*)-1-(1,2-dioxo-2-phenyl)ethyl-2-pyrrolidinecarboxylate;
- 1,7-diphenyl-4-heptyl      (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;
- 3-phenyl-1-propyl      (2*S*)-1-(3,3-dimethyl-1,2-dioxo-4-hydroxybutyl)-2-pyrrolidinecarboxylate;
- 20      3-phenyl-1-propyl      (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxamide;
- 1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylalanine ethyl ester;
- 1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-leucine ethyl ester;
- 25      1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylglycine ethyl ester;
- 1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-

phenylalanine phenyl ester;


1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylalanine benzyl ester;

1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-isoleucine ethyl ester; and

pharmaceutically acceptable salts, esters, and solvates thereof.

### FORMULA II

The pyrrolidine derivative may also be a compound of formula II




or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

20  $R_1$  is  $C_1$ - $C_8$  straight or branched chain alkyl,  $C_2$ - $C_8$  straight or branched chain alkenyl,  $C_3$ - $C_8$  cycloalkyl,  $C_5$ - $C_7$  cycloalkenyl or  $Ar_1$ , wherein said  $R_1$  is unsubstituted or substituted with one or more substituents independently selected from the group

25 consisting of  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_3$ - $C_8$  cycloalkyl,  $C_5$ - $C_7$  cycloalkenyl, hydroxy, and  $Ar_2$ ;

$Ar_1$  and  $Ar_2$  are independently selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl,

- 3-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl and phenyl, wherein said  $Ar_1$  is unsubstituted or substituted with one or more substituent(s) independently selected from the group consisting of hydrogen, halo, hydroxy, nitro, trifluoromethyl,  $C_1-C_6$  straight or branched chain alkyl,  $C_2-C_6$  straight or branched chain alkenyl,  $C_1-C_4$  alkoxy,  $C_2-C_4$  alkenyloxy, phenoxy, benzyloxy, and amino;
- 10           Z is  $C_1-C_6$  straight or branched chain alkyl, or  $C_2-C_6$  straight or branched chain alkenyl, wherein said Z is substituted with one or more substituent(s) independently selected from the group consisting of  $Ar_1$ ,  $C_3-C_8$  cycloalkyl, and  $C_1-C_6$  straight or branched chain alkyl or  $C_2-C_6$  straight or branched chain alkenyl substituted with  $C_3-C_8$  cycloalkyl; or Z is fragment
- 15



wherein:

- $R_3$  is  $C_1-C_6$  straight or branched chain alkyl which is unsubstituted or substituted with  $C_3-C_8$  cycloalkyl or  $Ar_1$ ;
- 25

$X_2$  is O or  $NR_5$ , wherein  $R_5$  is selected from the group consisting of hydrogen,  $C_1-C_6$  straight or branched chain alkyl, and  $C_2-C_6$  straight or branched

chain alkenyl; and

R<sub>4</sub> is selected from the group consisting of phenyl, benzyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl substituted with phenyl, and C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl substituted with phenyl.


In a preferred embodiment of formula II,  $R_1$  is selected from the group consisting of  $C_1$ - $C_9$ , straight or branched chain alkyl, 2-cyclohexyl, 4-cyclohexyl, 2-furanyl, 2-thienyl, 2-thiazolyl, and 4-hydroxybutyl.

In another preferred embodiment of formula II, Z and R<sub>1</sub> are lipophilic.

## 15

**FORMULA III**

The pyrrolidine derivative may also be a compound of formula III




III

25

or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

Z' is fragment



5

wherein:

R<sub>3</sub> is C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl or unsubstituted Ar<sub>1</sub>, wherein said alkyl is unsubstituted or substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl or Ar<sub>1</sub>;

10

X<sub>2</sub> is O or NR<sub>5</sub>, wherein R<sub>5</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, and C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl;

15

R<sub>4</sub> is selected from the group consisting of phenyl, benzyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl substituted with phenyl, and C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl substituted with phenyl; and

20


Ar<sub>1</sub> is as defined in formula II.

In a preferred embodiment of formula III, Z' is lipophilic.

#### FORMULA IV

25

Additionally, the pyrrolidine derivative may be a compound of formula IV



IV

5

wherein:

$R_1$  is  $C_1$ - $C_6$  straight or branched chain alkyl,  $C_2$ - $C_6$  straight or branched chain alkenyl,  $C_3$ - $C_6$  cycloalkyl or  $Ar_1$ , wherein said alkyl or alkenyl is unsubstituted or substituted with  $C_3$ - $C_6$  cycloalkyl or  $Ar_2$ ;

10

$Ar_1$  and  $Ar_2$  are independently selected from the group consisting of 2-furyl, 2-thienyl, and phenyl;

15

$X$  is selected from the group consisting of oxygen and sulfur;

$Y$  is oxygen;

20

$Z$  is  $C_1$ - $C_6$  straight or branched chain alkyl, or  $C_2$ - $C_6$  straight or branched chain alkenyl, wherein said  $Z$  is substituted with one or more substituent(s) independently selected from the group consisting of 2-furyl, 2-thienyl,  $C_3$ - $C_6$  cycloalkyl, pyridyl, and phenyl, each having one or more substituent(s) independently selected from the group consisting of hydrogen and  $C_1$ - $C_4$  alkoxy.

25

In a preferred embodiment of formula IV,  $Z$  and  $R_1$  are lipophilic.

In another preferred embodiment of formula IV, the compound is selected from the group consisting of:

dioxopentyl)-2-pyrrolidinecarboxylate;

10           3-(2-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

          3-(4-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

          3-phenyl-1-propyl       (2S)-1-(2-tert-butyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

15           3-phenyl-1-propyl (2S)-1-(2-cyclohexylethyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

          3-(3-pyridyl)-1-propyl (2S)-1-(2-cyclohexylethyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

20           3-(3-pyridyl)-1-propyl (2S)-1-(2-tert-butyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

          3,3-diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

          3-(3-pyridyl)-1-propyl (2S)-1-(2-cyclohexyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

25           3-(3-pyridyl)-1-propyl       (2S)-N-([2-thienyl]glyoxyl)pyrrolidinecarboxylate;

          3,3-diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-



dioxobutyl)-2-pyrrolidinecarboxylate;

3,3-diphenyl-1-propyl (2*S*)-1-cyclohexylglyoxyl-  
2-pyrrolidinecarboxylate;

3,3-diphenyl-1-propyl (2*S*)-1-(2-thienyl)glyoxyl-  
5 2-pyrrolidinecarboxylate; and

pharmaceutically acceptable salts, esters, and  
solvates thereof.

In a more preferred embodiment of formula IV, the  
compound is selected from the group consisting of:

10 3-(3-pyridyl)-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-  
dioxopentyl)-2-pyrrolidinecarboxylate;

3-(2-pyridyl)-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-  
dioxopentyl)-2-pyrrolidinecarboxylate;


3-(3-pyridyl)-1-propyl (2*S*)-1-(2-cyclohexyl-1,2-  
15 dioxoethyl)-2-pyrrolidinecarboxylate; and

pharmaceutically acceptable salts, esters, and  
solvates thereof.

In the most preferred embodiment of formula IV,  
the compound is 3-(3-pyridyl)-1-propyl (2*S*)-1-(3,3-  
20 dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate,  
and pharmaceutically acceptable salts, esters, and  
solvates thereof.

#### FORMULA V

25 Additionally, the pyrrolidine derivative may be  
a compound of formula V



V

5

or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

10

V is C, N, or S;

15

A and B, taken together with V and the carbon atom to which they are respectively attached, form a 5-7 membered saturated or unsaturated heterocyclic ring containing, in addition to V, one or more heteroatom(s) selected from the group consisting of O, S, SO, SO<sub>2</sub>, N, NH, and NR;

20

R is either C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>9</sub> straight or branched chain alkenyl, C<sub>3</sub>-C<sub>9</sub> cycloalkyl, C<sub>5</sub>-C<sub>9</sub> cycloalkenyl, or Ar<sub>1</sub>, wherein R is either unsubstituted or substituted with one or more substituent(s) independently selected from the group consisting of halo, haloalkyl, carbonyl, carboxy, hydroxy, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, thioalkyl, alkylthio, sulfhydryl, amino, alkylamino, aminoalkyl, aminocarboxyl, and Ar<sub>2</sub>;

25

R<sub>1</sub> is C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>9</sub>


straight or branched chain alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl or Ar<sub>1</sub>, wherein said R<sub>1</sub> is unsubstituted or substituted with one or more substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, hydroxy, and Ar<sub>2</sub>;

Ar<sub>1</sub> and Ar<sub>2</sub> are independently an alicyclic or aromatic, mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either unsubstituted or substituted with one or more substituent(s); wherein the individual ring size is 5-8 members; wherein said heterocyclic ring contains 1-6 heteroatom(s) independently selected from the group consisting of O, N, and S;

X is O, S, CH<sub>2</sub> or H<sub>2</sub>;

Y is O or NR<sub>2</sub>, wherein R<sub>2</sub> is hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; and

Z is C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, wherein said Z is substituted with one or more substituent(s) independently selected from the group consisting of Ar<sub>1</sub>, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl; or Z is fragment




C<sub>2</sub>, C<sub>3</sub> straight or branched chain alkenyl substituted  
with phenyl.

15           All the compounds of Formulas I-V possess  
asymmetric centers and thus can be produced as  
mixtures of stereoisomers or as individual R- and S-  
stereoisomers. The individual stereoisomers may be  
obtained by using an optically active starting  
20   material, by resolving a racemic or non-racemic  
mixture of an intermediate at some appropriate stage  
of the synthesis, or by resolving the compounds of  
Formulas I-V. It is understood that the compounds of  
Formulas I-V encompass individual stereoisomers as  
25   well as mixtures (racemic and non-racemic) of  
stereoisomers. Preferably, S-stereoisomers are used  
in the pharmaceutical compositions and methods of the  
present invention.

### Synthesis of Pyrrolidine Derivatives

The compounds of formulas I to V may be prepared by a variety of synthetic sequences that utilize established chemical transformations. The general pathway to the present compounds is described in Scheme I. N-glyoxylproline derivatives may be prepared by reacting L-proline methyl ester with methyl oxalyl chloride as shown in Scheme I. The resulting oxamates may be reacted with a variety of carbon nucleophiles to obtain intermediate compounds. These intermediates are then reacted with a variety of alcohols, amides, or protected amino acid residues to obtain the propyl esters and amides of the invention.


#### SCHEME I



reduced to the saturated alcohols by reaction with  
 excess lithium aluminum hydride, or sequentially by  
 10 reduction of the double bond by catalytic  
 hydrogenation and reduction of the saturated ester by  
 appropriate reducing agents. Alternatively, the  
 trans-cinnamates may be reduced to (E)-allylic  
 alcohols by the use of diisobutylaluminum hydride.

15

### SCHEME II



Longer chain alcohols may be prepared by  
 25 homologation of benzylic and higher aldehydes.  
 Alternatively, these aldehydes may be prepared by  
 conversion of the corresponding phenylacetic and  
 higher acids, and phenethyl and higher alcohols.

### Affinity for FKBP12

The compounds used in the inventive methods and pharmaceutical compositions have an affinity for the FK506 binding protein, particularly FKBP12. The inhibition of the prolyl peptidyl *cis-trans* isomerase activity of FKBP may be measured as an indicator of this affinity.

### K<sub>i</sub> Test Procedure

Inhibition of the peptidyl-prolyl isomerase (rotamase) activity of the compounds used in the inventive methods and pharmaceutical compositions can be evaluated by known methods described in the literature (Harding et al., *Nature*, 1989, 341:758-760; Holt et al. *J. Am. Chem. Soc.*, 115:9923-9938). These values are obtained as apparent K<sub>i</sub>'s and are presented for representative compounds in TABLE I.


The *cis-trans* isomerization of an alanine-proline bond in a model substrate, N-succinyl-Ala-Ala-Pro-Phe-p-nitroanilide, is monitored spectrophotometrically in a chymotrypsin-coupled assay, which releases para-nitroanilide from the *trans* form of the substrate. The inhibition of this reaction caused by the addition of different concentrations of inhibitor is determined, and the data is analyzed as a change in first-order rate constant as a function of inhibitor concentration to yield the apparent K<sub>i</sub> values.

In a plastic cuvette are added 950 mL of ice cold assay buffer (25 mM HEPES, pH 7.8, 100 mM NaCl), 10 mL of FKBP (2.5 mM in 10 mM Tris-Cl pH 7.5, 100 mM NaCl, 1 mM dithiothreitol), 25 mL of chymotrypsin (50 mg/mL in 1 mM HCl) and 10 mL of test compound at various concentrations in dimethyl sulfoxide. The reaction is initiated by the addition of 5 mL of substrate (succinyl-Ala-Phe-Pro-Phe-para-nitroanilide, 5 mg/mL in 2.35 mM LiCl in trifluoroethanol).

The absorbance at 390 nm versus time is monitored for 90 seconds using a spectrophotometer and the rate constants are determined from the absorbance versus time data files.

**TABLE 1**

*In Vitro* Test Results - Formulas I to V





**TABLE I***In Vitro* Test Results - Formulas I to V

|    | No. | Z                  | R <sub>1</sub>                              | K <sub>i</sub> |
|----|-----|--------------------|---|----------------|
| 5  | 1   | 1,1-dimethylpropyl | 3-phenylpropyl                              | 42             |
|    | 2   | "                  | 3-phenyl-prop-2-(E)-enyl                    | 125            |
|    | 3   | "                  | 3-(3,4,5-trimethoxy-phenyl)propyl           | 200            |
| 10 | 4   | "                  | 3-(3,4,5-trimethoxy-phenyl)-prop-2-(E)-enyl | 65             |
| 15 | 5   | "                  | 3-(4,5-methylenedioxy)-phenylpropyl         | 170            |
|    | 6   | "                  | 3-(4,5-methylenedioxy)phenylprop-2-(E)-enyl | 160            |
| 20 | 7   | "                  | 3-cyclohexylpropyl                          | 200            |
|    | 8   | "                  | 3-cyclohexylprop-2-(E)-enyl                 | 600            |
|    | 9   | "                  | (1 <i>R</i> )-1,3-diphenyl-1-propyl         | 52             |
| 25 | 10  | 2-furanyl          | 3-phenylpropyl                              | 4000           |
|    | 11  | 2-thienyl          | "   | 92             |
| 30 | 12  | 2-thiazolyl        | "   | 100            |
|    | 13  | phenyl             | "   | 1970           |
| 35 | 14  | 1,1-dimethylpropyl | 3-(2,5-dimethoxy)phenylpropyl               | 250            |
|    | 15  | "                  | 3-(2,5-dimethoxy)phenylprop-2-(E)-enyl      | 450            |
| 40 | 16  | "                  | 2-(3,4,5-trimethoxyphenyl)ethyl             | 120            |
|    | 17  | "                  | 3-(3-pyridyl)propyl                         | 5              |
|    | 18  | "                  | 3-(2-pyridyl)propyl                         | 195            |

In Vitro Test Results - Formulas I to V

|    | No. | Z                  | R <sub>1</sub>      | K <sub>i</sub> |
|----|-----|--------------------|---------------------|----------------|
| 5  |     |                    |                     |                |
|    | 19  | "                  | 3-(4-pyridyl)propyl | 23             |
| 10 | 20  | cyclohexyl         | 3-phenylpropyl      | 82             |
|    | 21  | <i>tert</i> -butyl | "                   | 95             |
| 15 | 22  | cyclohexylethyl    | "                   | 1025           |
|    | 23  | cyclohexylethyl    | 3-(3-pyridyl)propyl | 1400           |
|    | 24  | <i>tert</i> -butyl | 3-(3-pyridyl)propyl | 3              |
| 20 | 25  | 1,1-dimethylpropyl | 3,3-diphenylpropyl  | 5              |
|    | 26  | cyclohexyl         | 3-(3-pyridyl)propyl | 9              |
| 25 | 27  | 2-thienyl          | 3-(3-pyridyl)propyl | 1000           |
|    | 28  | <i>tert</i> -butyl | 3,3-diphenylpropyl  | 5              |
|    | 29  | cyclohexyl         | "                   | 20             |
| 30 | 30  | 2-thienyl          | "                   | 150            |

Route of Administration

35           To effectively treat alopecia or promote hair growth, the compounds used in the inventive methods and pharmaceutical compositions must readily affect the targeted areas. For these purposes, the compounds are preferably administered topically to the skin.

40           For topical application to the skin, the compounds can be formulated into suitable ointments containing the compounds suspended or dissolved in,

for example, mixtures with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water.

- 5 Alternatively, the compounds can be formulated into suitable lotions or creams containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl ester
- 10 wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

Dosage levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred levels of about 0.1 mg to about 1,000 mg.

10 The specific dose level for any particular patient will vary depending upon a variety of factors, including the activity of the specific compound employed; the age, body weight, general health, sex and diet of the patient; the time of administration;  
15 the rate of excretion; drug combination; the severity of the particular disease being treated; and the form of administration. Typically, *in vitro* dosage-effect results provide useful guidance on the proper doses for patient administration. Studies in animal models  
20 are also helpful. The considerations for determining the proper dose levels are well known in the art.

The compounds can be administered with other hair revitalizing agents. Specific dose levels for the other hair revitalizing agents will depend upon the  
25 factors previously stated and the effectiveness of the drug combination.

EXAMPLES

The following examples are illustrative of the present invention and are not intended to be limitations thereon. Unless otherwise indicated, all percentages are based upon 100% by weight of the final composition.

EXAMPLE 1

Synthesis of 3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate (1)  
Methyl (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate

A solution of L-proline methyl ester hydrochloride (3.08 g; 18.60 mmol) in dry methylene chloride was cooled to 0°C and treated with triethylamine (3.92 g; 38.74 mmol; 2.1 eq). After stirring the formed slurry under a nitrogen atmosphere for 15 min, a solution of methyl oxalyl chloride (3.20 g; 26.12 mmol) in methylene chloride (45 mL) was added dropwise. The resulting mixture was stirred at 0°C for 1.5 hour. After filtering to remove solids, the organic phase was washed with water, dried over MgSO<sub>4</sub> and concentrated. The crude residue was purified on a silica gel column, eluting with 50% ethyl acetate in hexane, to obtain 3.52 g (88%) of the product as a reddish oil. Mixture of cis-trans amide rotamers; data for trans rotamer given. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 1.93 (dm, 2H); 2.17 (m, 2H); 3.62 (m, 2H); 3.71 (s, 3H);

5 methoxyethyl)-2-pyrrolidinecarboxylate (2.35 g; 10.90  
mmol) in 30 mL of tetrahydrofuran (THF) was cooled to  
-78°C and treated with 14.2 mL of a 1.0 M solution of  
1,1-dimethylpropylmagnesium chloride in THF. After  
10 stirring the resulting homogeneous mixture at -78°C for  
three hours, the mixture was poured into saturated  
ammonium chloride (100 mL) and extracted into ethyl  
acetate. The organic phase was washed with water,  
dried, and concentrated, and the crude material  
obtained upon removal of the solvent was purified on  
15 a silica gel column, eluting with 25% ethyl acetate in  
hexane, to obtain 2.10 g (75%) of the oxamate as a  
colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 0.88 (t, 3H); 1.22,  
1.26 (s, 3H each); 1.75 (dm, 2H); 1.87-2.10 (m, 3H);  
2.23 (m, 1H); 3.54 (m, 2H); 3.76 (s, 3H); 4.52 (dm,  
20 1H, J = 8.4, 3.4).

Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-  
pyrrolidinecarboxylic acid

A mixture of methyl (2S)-1-(1,2-dioxo-3,3-  
dimethylpentyl)-2-pyrrolidinecarboxylate (2.10 g; 8.23  
25 mmol), 1 N LiOH (15 mL), and methanol (50 mL) was  
stirred at 0°C for 30 minutes and at room temperature  
overnight. The mixture was acidified to pH 1 with 1 N  
HCl, diluted with water, and extracted into 100 mL of

methylene chloride. The organic extract was washed with brine and concentrated to deliver 1.73 g (87%) of snow-white solid which did not require further purification.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): d 0.87 (t, 3H); 1.22, 1.25 (s, 3H each); 1.77 (dm, 2H); 2.02 (m, 2H); 2.17 (m, 1H); 2.25 (m, 1H); 3.53 (dd, 2H,  $J = 10.4, 7.3$ ); 4.55 (dd, 1H,  $J = 8.6, 4.1$ ).

3-Phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate (1)

10           A mixture of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid (600 mg; 2.49 mmol), 3-phenyl-1-propanol (508 mg; 3.73 mmol), dicyclohexylcarbodiimide (822 mg; 3.98 mmol), camphorsulfonic acid (190 mg; 0.8 mmol) and 4-  
15           dimethylaminopyridine (100 mg; 0.8 mmol) in methylene chloride (20 mL) was stirred overnight under a nitrogen atmosphere. The reaction mixture was filtered through Celite to remove solids and concentrated in vacuo, and the crude material was  
20           purified on a flash column (25% ethyl acetate in hexane) to obtain 720 mg (80%) of Example 1 as a colorless oil.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ): d 0.84 (t, 3H); 1.19 (s, 3H); 1.23 (s, 3H); 1.70 (dm, 2H); 1.98 (m, 5H); 2.22 (m, 1H); 2.64 (m, 2H); 3.47 (m, 2H); 4.14 (m, 2H); 4.51 (d, 1H); 7.16 (m, 3H); 7.26 (m, 2H).  
25

Example 2

The method of Example 1 was utilized to prepare the following illustrative compounds.

5      Compound 2: 3-phenyl-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, 80%. <sup>1</sup>H NMR (360 Mhz, CDCl<sub>3</sub>): d 0.86 (t, 3H); 1.21 (s, 3H); 1.25 (s, 3H); 1.54-2.10 (m, 5H); 2.10-2.37 (m, 1H); 3.52-3.55 (m, 2H); 4.56 (dd, 1H, J = 3.8, 8.9); 4.78-4.83 (m, 2H); 6.27 (m, 1H); 6.67 (dd, 1H, J = 15.9); 7.13-7.50 (m, 5H).

15      Compound 3: 3-(3,4,5-trimethoxyphenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, 61%. <sup>1</sup>H NMR (CDCl<sub>3</sub>): d 0.84 (t, 3H); 1.15 (s, 3H); 1.24 (s, 3H); 1.71 (dm, 2H); 1.98 (m, 5H); 2.24 (m, 1H); 2.63 (m, 2H); 3.51 (t, 2H); 3.79 (s, 3H); 3.83 (s, 3H); 4.14 (m, 2H); 4.52 (m, 1H); 6.36 (s, 2H).

20      Compound 4: 3-(3,4,5-trimethoxyphenyl)-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidine carboxylate, 66%. <sup>1</sup>H NMR (CDCl<sub>3</sub>): d 0.85 (t, 3H); 1.22 (s, 3H); 1.25 (s, 3H); 1.50-2.11 (m, 5H); 2.11-2.40 (m, 1H); 3.55 (m, 2H); 3.85 (s, 3H); 3.88 (s, 6H); 4.56 (dd, 1H); 4.81 (m, 2H); 6.22 (m, 1H); 6.58 (d, 1H, J = 16); 6.63 (s, 2H).



- Compound 5: 3-(4,5-methylenedioxyphenyl)-1-propyl  
(2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidine-  
carboxylate, 82%. <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>): d 0.86 (t,  
3H); 1.22 (s, 3H); 1.25 (s, 3H); 1.60-2.10 (m, 5H);  
5 3.36-3.79 (m, 2H); 4.53 (dd, 1H, *J* = 3.8, 8.6); 4.61-  
4.89 (m, 2H); 5.96 (s, 2H); 6.10 (m, 1H); 6.57 (dd,  
1H, *J* = 6.2, 15.8); 6.75 (d, 1H, *J* = 8.0); 6.83 (dd,  
1H, *J* = 1.3, 8.0); 6.93 (s, 1H).
- 10 Compound 6: 3-(4,5-methylenedioxyphenyl)-1-prop-2-  
(*E*)-enyl (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-  
pyrrolidinecarboxylate, 82%. <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>):  
d 0.86 (t, 3H); 1.22 (s, 3H); 1.25 (s, 3H); 1.60-2.10  
(m, 5H); 2.10-2.39 (m, 1H); 3.36-3.79 (m, 2H); 4.53  
15 (dd, 1H, *J* = 3.8, 8.6); 4.61-4.89 (m, 2H); 5.96 (s,  
2H); 6.10 (m, 1H); 6.57 (dd, 1H, *J* = 6.2, 15.8); 6.75  
(d, 1H, *J* = 8.0); 6.83 (dd, 1H, *J* = 1.3, 8.0); 6.93  
(s, 1H).
- 20 Compound 8: 3-cyclohexyl-1-prop-2-(*E*)-enyl (2*S*)-1-  
(3,3-dimethyl-1,2-dioxopentyl)-2-  
pyrrolidinecarboxylate, 92%. <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>):  
d 0.86 (t, 3H); 1.13-1.40 (m + 2 singlets, 9H total);  
1.50-1.87 (m, 8H); 1.87-2.44 (m, 6H); 3.34-3.82 (m,  
25 2H); 4.40-4.76 (m, 3H); 5.35-5.60 (m, 1H); 5.60-5.82  
(dd, 1H, *J* = 6.5, 16).

Compound 9: (1*R*)-1,3-Diphenyl-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, 90%. <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>): d 0.85 (t, 3H); 1.20 (s, 3H); 1.23 (s, 3H); 1.49-2.39 (m, 7H); 2.46-2.86 (m, 2H); 3.25-3.80 (m, 2H); 4.42-4.82 (m, 1H); 5.82 (td, 1H, *J* = 1.8, 6.7); 7.05-7.21 (m, 3H); 7.21-7.46 (m, 7H).

Compound 10: 3-phenyl-1-propyl (2*S*)-1-(1,2-dioxo-2-[2-furanyl])ethyl-2-pyrrolidinecarboxylate, 99%. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): d 1.66-2.41 (m, 6H); 2.72 (t, 2H, *J* = 7.5); 3.75 (m, 2H); 4.21 (m, 2H); 4.61 (m, 1H); 6.58 (m, 1H); 7.16-7.29 (m, 5H); 7.73 (m, 2H).

Compound 11: 3-phenyl-1-propyl (2*S*)-1-(1,2-dioxo-2-[2-thienyl])ethyl-2-pyrrolidinecarboxylate, 81%. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): d 1.88-2.41 (m, 6H); 2.72 (dm, 2H); 3.72 (m, 2H); 4.05 (m, 1H); 4.22 (m, 1H); 4.64 (m, 1H); 7.13-7.29 (m, 6H); 7.75 (dm, 1H); 8.05 (m, 1H).

Compound 13: 3-phenyl-1-propyl (2*S*)-1-(1,2-dioxo-2-phenyl)ethyl-2-pyrrolidinecarboxylate, 99%. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): d 1.97-2.32 (m, 6H); 2.74 (t, 2H, *J* = 7.5); 3.57 (m, 2H); 4.24 (m, 2H); 4.67 (m, 1H); 6.95-7.28 (m, 5H); 7.51-7.64 (m, 3H); 8.03-8.09 (m, 2H).

Compound 14: 3-(2,5-dimethoxyphenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidine-carboxylate, 99%. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): d 0.87 (t, 3H); 1.22 (s, 3H); 1.26 (s, 3H); 1.69 (m, 2H); 1.96 (m, 5H); 2.24 (m, 1H); 2.68 (m, 2H); 3.55 (m, 2H); 3.75 (s, 3H); 3.77 (s, 3H); 4.17 (m, 2H); 4.53 (d, 1H); 6.72 (m, 3H).

Compound 15: 3-(2,5-dimethoxyphenyl)-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidine-carboxylate, 99%. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): d 0.87 (t, 3H); 1.22 (s, 3H); 1.26 (s, 3H); 1.67 (m, 2H); 1.78 (m, 1H); 2.07 (m, 2H); 2.26 (m, 1H); 3.52 (m, 2H); 3.78 (s, 3H); 3.80 (s, 3H); 4.54 (m, 1H); 4.81 (m, 2H); 6.29 (dt, 1H, J = 15.9); 6.98 (s, 1H).

Compound 16: 2-(3,4,5-trimethoxyphenyl)-1-ethyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidine-carboxylate, 97%. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): d 0.84 (t, 3H); 1.15 (s, 3H); 1.24 (s, 3H); 1.71 (dm, 2H); 1.98 (m, 5H); 2.24 (m, 1H); 2.63 (m, 2H); 3.51 (t, 2H); 3.79 (s, 3H); 3.83 (s, 3H); 4.14 (m, 2H); 4.52 (m, 1H); 6.36 (s, 2H).

25

Compound 17: 3-(3-Pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, 80%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): d 0.85 (t, 3H); 1.23,

10 7.3), 3.52 (m, 2H), 4.20 (m, 2H); 4.51 (m, 1H); 7.09-  
7.19 (m, 2H); 7.59 (m, 1H); 8.53 (d, 1H,  $J = 4.9$ ).

Compound 19: 3-(4-Pyridyl)-1-propyl (2S)-1-(3,3-  
dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate,  
15 91%.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz): d 6.92-6.80 (m, 4H);  
6.28 (m, 1H); 5.25 (d, 1H,  $J = 5.7$ ); 4.12 (m, 1H);  
4.08 (s, 3H); 3.79 (s, 3H); 3.30 (m, 2H); 2.33 (m,  
1H); 1.85-1.22 (m, 7H); 1.25 (s, 3H); 1.23 (s, 3H);  
0.89 (t, 3H,  $J = 7.5$ ).

20 Compound 20: 3-phenyl-1-propyl (2S)-1-(2-cyclohexyl-  
1,2-dioxoethyl)-2-pyrrolidinecarboxylate, 91%.  $^1\text{H}$  NMR  
( $\text{CDCl}_3$ , 300 MHz): d 1.09-1.33 (m, 5H); 1.62-2.33 (m,  
12H); 2.69 (t, 2H,  $J = 7.5$ ); 3.15 (dm, 1H); 3.68 (m,  
2H); 4.16 (m, 2H); 4.53, 4.84 (d, 1H total); 7.19 (m,  
25 3H); 7.29 (m, 2H).

Compound 21: 3-phenyl-1-propyl (2S)-1-(2-tert-butyl-  
1,2-dioxoethyl)-2-pyrrolidinecarboxylate, 92%.  $^1\text{H}$  NMR

(CDCl<sub>3</sub>, 300 MHz): d 1.29 (s, 9H); 1.94-2.03 (m, 5H); 2.21 (m, 1H); 2.69 (m, 2H); 3.50-3.52 (m, 2H); 4.16 (m, 2H); 4.53 (m, 1H); 7.19 (m, 3H); 7.30 (m, 2H).

5      Compound 22: 3-phenyl-1-propyl (2*S*)-1-(2-cyclohexylethyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate, 97%.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): d 0.88 (m, 2H); 1.16 (m, 4H); 1.43-1.51 (m, 2H); 1.67 (m, 5H); 1.94-2.01 (m, 6H); 2.66-2.87 (m, 4H); 3.62-3.77 (m, 2H); 4.15 (m, 2H);  
 10      4.86 (m, 1H); 7.17-7.32 (m, 5H).

Compound 23: 3-(3-pyridyl)-1-propyl (2*S*)-1-(2-cyclohexylethyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate, 70%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): d 0.87 (m, 2H); 1.16 (m, 4H); 1.49 (m, 2H); 1.68 (m, 4H); 1.95-2.32 (m, 7H); 2.71 (m, 2H); 2.85 (m, 2H); 3.63-3.78 (m, 2H); 4.19 (m, 2H); 5.30 (m, 1H); 7.23 (m, 1H); 7.53 (m, 1H); 8.46 (m, 2H).

20      Compound 24: 3-(3-pyridyl)-1-propyl (2*S*)-1-(2-tert-butyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate, 83%.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): d 1.29 (s, 9H); 1.95-2.04 (m, 5H); 2.31 (m, 1H); 2.72 (t, 2H, J = 7.5); 3.52 (m, 2H); 4.18 (m, 2H); 4.52 (m, 1H); 7.19-7.25 (m, 1H);  
 25      7.53 (m, 1H); 8.46 (m, 2H).

Compound 25: 3,3-diphenyl-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate,

99%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): d 0.85 (t, 3H); 1.21, 1.26 (s, 3H each); 1.68-2.04 (m, 5H); 2.31 (m, 1H); 2.40 (m, 2H); 3.51 (m, 2H); 4.08 (m, 3H); 4.52 (m, 1H); 7.18-7.31 (m, 10H).

5

Compound 26: 3-(3-pyridyl)-1-propyl (2S)-1-(2-cyclohexyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate, 88%.

<sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): d 1.24-1.28 (m, 5H); 1.88-2.35 (m, 11H); 2.72 (t, 2H, J = 7.5); 3.00-3.33 (dm, 1H); 3.69 (m, 2H); 4.19 (m, 2H); 4.55 (m, 1H); 7.20-7.24 (m, 1H); 7.53 (m, 1H); 8.47 (m, 2H).

10

Compound 27: 3-(3-Pyridyl)-1-propyl (2S)-N-([2-thienyl] glyoxyl)pyrrolidinecarboxylate, 49%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): d 1.81-2.39 (m, 6H); 2.72 (dm, 2H); 3.73 (m, 2H); 4.21 (m, 2H); 4.95 (m, 1H); 7.19 (m, 2H); 7.61 (m, 1H); 7.80 (d, 1H); 8.04 (d, 1H); 8.46 (m, 2H).

15

Compound 28: 3,3-Diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxobutyl)-2-pyrrolidinecarboxylate, 99%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): d 1.27 (s, 9H); 1.96 (m, 2H); 2.44 (m, 4H); 3.49 (m, 1H); 3.64 (m, 1H); 4.08 (m, 4H); 4.53 (dd, 1H); 7.24 (m, 10H).

20

25

Compound 29: 3,3-Diphenyl-1-propyl (2S)-1-cyclohexyl glyoxyl-2-pyrrolidinecarboxylate, 91%. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): d 1.32 (m, 6H); 1.54-2.41 (m, 10H); 3.20

(dm, 1H); 3.69 (m, 2H); 4.12 (m, 4H); 4.52 (d, 1H);  
7.28 (m, 10H).

Compound 30: 3,3-Diphenyl-1-propyl (2*S*)-1-(2-thienyl)  
5 glyoxyl-2-pyrrolidinecarboxylate, 75%. <sup>1</sup>H NMR (CDCl<sub>3</sub>,  
300 MHz): d 2.04 (m, 3H); 2.26 (m, 2H); 2.48 (m, 1H);  
3.70 (m, 2H); 3.82-4.18 (m, 3H total); 4.64 (m, 1H);  
7.25 (m, 11H); 7.76 (dd, 1H); 8.03 (m, 1H).

10

### Example 3

General procedure for the synthesis of acrylic  
esters, exemplified for methyl (3,3,5-trimethoxy)-  
*trans*-cinnamate.

A solution of 3,4,5-trimethoxybenzaldehyde (5.0  
15 g; 25.48 mmol) and methyl (triphenyl-  
phosphoranylidene)acetate (10.0 g; 29.91 mmol) in  
tetrahydrofuran (250 mL) was refluxed overnight.  
After cooling, the reaction mixture was diluted with  
200 mL of ethyl acetate and washed with 2 x 200 mL of  
20 water, dried, and concentrated in vacuo. The crude  
residue was chromatographed on a silica gel column,  
eluting with 25% ethyl acetate in hexane, to obtain  
5.63 g (88%) of the cinnamate as a white crystalline  
solid. <sup>1</sup>H NMR (300 Mhz; CDCl<sub>3</sub>): d 3.78 (s, 3H); 3.85  
25 (s, 6H); 6.32 (d, 1H, *J* = 16); 6.72 (s, 2H); 7.59 (d,  
1H, *J* = 16).

Example 4

General procedure for the synthesis of saturated alcohols from acrylic esters, exemplified for (3,4,5-trimethoxy) phenylpropanol.

5           A solution of methyl (3,3,5-trimethoxy)-*trans*-cinnamate (1.81 g; 7.17 mmol) in tetrahydrofuran (30 mL) was added in a dropwise manner to a solution of lithium aluminum hydride (14 mmol) in THF (35 mL), with stirring and under an argon atmosphere. After  
10 the addition was complete, the mixture was heated to 75°C for 4 hours. After cooling, it was quenched by the careful addition of 15 mL of 2 N NaOH followed by 50 mL of water. The resulting mixture was filtered through Celite to remove solids, and the filter cake  
15 was washed with ethyl acetate. The combined organic fractions were washed with water, dried, concentrated in vacuo, and purified on a silica gel column, eluting with ethyl acetate to obtain 0.86 g (53%) of the alcohol as a clear oil. <sup>1</sup>H NMR (300 Mhz; CDCl<sub>3</sub>): d  
20 1.23 (br, 1H); 1.87 (m, 2H); 2.61 (t, 2H, J = 7.1); 3.66 (t, 2H); 3.80 (s, 3H); 3.83 (s, 6H); 6.40 (s, 2H).

Example 5

25           General procedure for the synthesis of *trans*-allylic alcohols from acrylic esters, exemplified for (3,4,5-trimethoxy)phenylprop-2-(E)-enol.

A solution of methyl (3,3,5-trimethoxy)-*trans*-



cinamate (1.35 g; 5.35 mmol) in toluene (25 mL) was cooled to  $-10^{\circ}\text{C}$  and treated with a solution of diisobutylaluminum hydride in toluene (11.25 mL of a 1.0 M solution; 11.25 mmol). The reaction mixture was stirred for 3 hours at  $0^{\circ}\text{C}$  and then quenched with 3 mL of methanol followed by 1 N HCl until the pH was 1. The reaction mixture was extracted into ethyl acetate and the organic phase was washed with water, dried and concentrated. Purification on a silica gel column eluting with 25% ethyl acetate in hexane furnished 0.96 g (80%) of a thick oil.  $^1\text{H}$  NMR (360 Mhz;  $\text{CDCl}_3$ ):  $\delta$  3.85 (s, 3H); 3.87 (s, 6H); 4.32 (d, 2H,  $J = 5.6$ ); 6.29 (dt, 1H,  $J = 15.8, 5.7$ ), 6.54 (d, 1H,  $J = 15.8$ ); 6.61 (s, 2H).

15

### Example 6

#### In Vivo Hair Generation Tests With C57 Black 6 Mice

Experiment A: C57 black 6 mice were used to demonstrate the hair revitalizing properties of a low molecular weight, small molecule non-immunosuppressive neuroimmunophilin FKBP ligand, GPI 1046, a pyrrolidine derivative. Referring now to FIGS. 1 and 2 of the drawings, C57 black 6 mice, approximately 7 weeks old, had an area of about 2 inches by 2 inches on their hindquarters shaved to remove all existing hair. Care was taken not to nick or cause abrasion to the underlaying dermal layers. The animals were in anagen growth phase, as indicated by the pinkish color of the

20

25

(FIG. 3) or 30  $\mu$ M GPI 1046 (FIG. 4) dissolved in the  
5 vehicle. The animals were treated with vehicle or GPI  
1046 every 48 hours (3 applications total over the  
course of 5 days) and the hair growth was allowed to  
proceed for 6 weeks. Hair growth was quantitated by  
the percent of shaved area covered by new hair growth  
10 during this time period.

FIG. 2 shows that animals treated with vehicle  
exhibited only a small amount of hair growth in  
patches or tufts, with less than 3% of the shaved area  
covered with new growth. In contrast, FIG. 3 shows  
15 that animals treated with 10  $\mu$ M GPI 1046 exhibited  
dramatic hair growth, covering greater than 90% of the  
shaved area in all animals. Further, FIG. 4 shows  
that mice treated with 30  $\mu$ M GPI 1046 exhibited  
essentially complete hair regrowth and their shaved  
20 areas were indistinguishable from unshaven C57 black  
6 mice.

Experiment B: C57 Black 6 mice were used to  
demonstrate the hair revitalizing properties of  
various low molecular weight, small molecule,  
25 non-immunosuppressive neuroimmunophilin FKBP ligands,  
including GPI 1046. C57 Black 6 mice, 55 to 75 days  
old, had an area of about 2 inches by 2 inches on  
their hindquarters shaved to remove all existing hair.

Care was taken not to nick or cause abrasion to the underlying dermal layers. The animals were in anagen growth phase when shaved. Five animals per group were treated by topical administration with a vehicle, FK506, or one of the low molecular weight, small molecule, non-immunosuppressive neuroimmunophilin FKBP ligands (GPI 1605, 1046, 1312, 1572, 1389, 1511, and 1234) at a concentration of one micromole per milliliter to the shaved area. The animals were treated three times per week, and hair growth was evaluated 14 days after initiation of treatment. Hair growth was quantitated by the percent of shaved area covered by new hair growth, as scored by a blinded observer, on a scale of 0 (no growth) to five (complete hair regrowth in shaved area).

Figure 5 shows that after 14 days, the animals treated with vehicle exhibited the beginning of growth in small tufts. In contrast, animals treated with one of the low molecular weight, small molecule, non-immunosuppressive neuroimmunophilin FKBP ligands, including GPI 1046, exhibited dramatic hair growth.

Example 7

A lotion comprising the following composition may be prepared.

|    |   |      |
|----|---|------|
| 5  |   | (%)  |
|    | 95% Ethanol   | 80.0 |
|    | a pyrrolidine derivative as defined above               | 10.0 |
|    | $\alpha$ -Tocopherol acetate                            | 0.01 |
| 10 | Ethylene oxide (40 mole) adducts of hardened castor oil | 0.5  |
|    | purified water  | 9.0  |
|    | perfume and dye   | q.s. |

15            Into 95% ethanol are added a pyrrolidine derivative,  $\alpha$ -tocopherol acetate, ethylene oxide (40 mole) adducts of hardened castor oil, perfume and a dye. The resulting mixture is stirred and dissolved, and purified water is added to the mixture to obtain  
20            a transparent liquid lotion.

             5 ml of the lotion may be applied once or twice per day to a site having marked baldness or alopecia.

Example 8

A lotion comprising the following composition shown may be prepared.

|    |   |       |
|----|---|-------|
| 5  |   | (%)   |
|    | 95% Ethanol   | 80.0  |
|    | a pyrrolidine derivative as defined above               | 0.005 |
|    | Hinokitol   | 0.01  |
| 10 | Ethylene oxide (40 mole) adducts of hardened castor oil | 0.5   |
|    | Purified water  | 19.0  |
|    | Perfume and dye   | q.s.  |

15 Into 95% ethanol are added a pyrrolidine derivative, hinokitol, ethylene oxide (40 mole) adducts of hardened castor oil, perfume, and a dye. The resulting mixture is stirred, and purified water is added to the mixture to obtain a transparent liquid lotion.

20 The lotion may be applied by spraying once to 4 times per day to a site having marked baldness or alopecia.

Example 9

An emulsion may be prepared from A phase and B phase having the following compositions.

|    |   |      |
|----|---|------|
| 5  | (A phase)                                 | (%)  |
|    | Whale wax                                 | 0.5  |
|    | Cetanol                                   | 2.0  |
|    | Petrolatum                                | 5.0  |
|    | Squalane                                  | 10.0 |
| 10 | Polyoxyethylene (10 mole) monostearate    | 2.0  |
|    | Sorbitan monooleate                       | 1.0  |
|    | a pyrrolidine derivative as defined above | 0.01 |
|    | (B phase)                                 | (%)  |
|    | Glycerine                                 | 10.0 |
| 15 | Purified water                            | 69.0 |
|    | Perfume, dye, and preservative            | q.s. |

The A phase and the B phase are respectively heated and melted and maintained at 80°C. Both phases are then mixed and cooled under stirring to normal temperature to obtain an emulsion.

The emulsion may be applied by spraying once to four times per day to a site having marked baldness or alopecia.

Example 10

A cream may be prepared from A phase and B phase having the following compositions.

|    |  |        |
|----|--|--------|
| 5  | (A Phase)                                      | (%)    |
|    | Fluid paraffin                                 | 5.0    |
|    | Cetostearyl alcohol                            | 5.5    |
|    | Petrolatum                                     | 5.5    |
|    | Glycerine monostearate                         | 33.0   |
| 10 | Polyoxyethylene (20 mole) 2-octyldodecyl ether | 3.0    |
|    | Propylparaben                                  | 0.3    |
|    | (B Phase)                                      | (%)    |
|    | a pyrrolidine derivative as defined above      | 0.8    |
| 15 | Glycerine                                      | 7.0    |
|    | Dipropylene glycol                             | 20.0   |
|    | Polyethylene glycol 4000                       | 5.0    |
|    | Sodium Hexametaphosphate                       | 0.005  |
| 20 | Purified water                                 | 44.895 |

The A phase is heated and melted, and maintained at 70°C. The B phase is added into the A phase and the mixture is stirred to obtain an emulsion. The emulsion is then cooled to obtain a cream.

25 The cream may be applied once to 4 times per day to a site having marked baldness or alopecia.

Example 11

A liquid comprising the following composition may be prepared.

|    |   |       |
|----|---|-------|
| 5  |   | (%)   |
|    | Polyoxyethylene butyl ether   | 20.0  |
|    | Ethanol   | 50.0  |
|    | a pyrrolidine derivative as defined above                                       | 0.001 |
|    | Propylene glycol  | 5.0   |
| 10 | Polyoxyethylene hardened castor oil derivative (ethylene oxide 80 mole adducts) | 0.4   |
|    | Perfume   | q.s.  |
|    | Purified water  | q.s.  |

15            Into ethanol are added polyoxypropylene butyl ether, propylene glycol, polyoxyethylene hardened castor oil, a pyrrolidine derivative, and perfume. The resulting mixture is stirred, and purified water is added to the mixture to obtain a liquid.

20            The liquid may be applied once to 4 times per day to a site having marked baldness or alopecia.



Example 12

A shampoo comprising the following composition may be prepared.

|    |   |      |
|----|---|------|
| 5  |   | (%)  |
|    | Sodium laurylsulfate                      | 5.0  |
|    | Triethanolamine laurylsulfate             | 5.0  |
|    | Betaine lauryldimethylaminoacetate        | 6.0  |
|    | Ethylene glycol distearate                | 2.0  |
| 10 | Polyethylene glycol                       | 5.0  |
|    | a pyrrolidine derivative as defined above | 5.0  |
|    | Ethanol                                   | 2.0  |
|    | Perfume                                   | 0.3  |
| 15 | Purified water                            | 69.7 |

Into 69.7 of purified water are added 5.0 g of sodium laurylsulfate, 5.0 g of triethanolamine laurylsulfate, 6.0 g of betaine lauryldimethylaminoacetate. Then a mixture obtained by adding 5.0 g of a pyrrolidine derivative, 5.0 g of polyethylene glycol, and 2.0 g of ethylene glycol distearate to 2.0 g of ethanol, followed by stirring, and 0.3 g of perfume are successively added. The resulting mixture is heated and subsequently cooled to obtain a shampoo.

The shampoo may be used on the scalp once or twice per day.

Example 13

A patient is suffering from alopecia senilis. A pyrrolidine derivative as identified above, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 14

A patient is suffering from male pattern alopecia. A pyrrolidine derivative as identified above, or a pharmaceutical composition comprising the same may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 15

A patient is suffering from alopecia areata. A pyrrolidine derivative as identified above, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 16

A patient is suffering from hair loss caused by skin lesions. A pyrrolidine derivative as identified above, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 17

A patient is suffering from hair loss caused by tumors. A pyrrolidine derivative as identified above, or a pharmaceutical composition comprising the same,  
5 may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 18

A patient is suffering from hair loss caused by  
10 a systematic disorder, such as a nutritional disorder or an internal secretion disorder. A pyrrolidine derivative as identified above, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to  
15 occur following treatment.

Example 19

A patient is suffering from hair loss caused by chemotherapy. A pyrrolidine derivative as identified  
20 above, or a pharmaceutical composition comprising the same, may be administered to the patient. Increased hair growth is expected to occur following treatment.

Example 20

25 A patient is suffering from hair loss caused by radiation. A pyrrolidine derivative as identified above, or a pharmaceutical composition comprising the

same may, be administered to the patient. Increased hair growth is expected to occur following treatment.

5       The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modifications are intended to be included within the scope of the following claims.

## WE CLAIM:

1. A method for treating alopecia or promoting hair growth in an animal, which comprises administering to said animal an effective amount of a pyrrolidine derivative.

2. The method of claim 1, wherein the pyrrolidine derivative is non-immunosuppressive.

10


3. The method of claim 1, wherein the pyrrolidine derivative has an affinity for an FKBP-type immunophilin.

15

4. The method of claim 3, wherein the FKBP-type immunophilin is FKBP-12.

5. The method of claim 1, wherein the pyrrolidine derivative is a compound of formula I

20



I

25

or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

$R_1$  is  $C_1$ - $C_9$  straight or branched chain alkyl,  $C_2$ - $C_9$ ,


straight or branched chain alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl or Ar<sub>1</sub>, wherein said R<sub>1</sub> is unsubstituted or substituted with one or more substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, hydroxy, and Ar<sub>2</sub>;

Ar<sub>1</sub> and Ar<sub>2</sub> are independently selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl and phenyl, wherein said Ar<sub>1</sub> is unsubstituted or substituted with one or more substituent(s) independently selected from the group consisting of hydrogen, halo, hydroxy, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, S, CH<sub>2</sub> or H<sub>2</sub>;

Y is O or NR<sub>2</sub>, wherein R<sub>2</sub> is hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; and

Z is C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, wherein said Z is substituted with one or more substituent(s) independently selected from the group consisting of Ar<sub>1</sub>, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl; or Z is fragment



5

wherein:

$\text{R}_3$  is  $\text{C}_1$ - $\text{C}_9$  straight or branched chain alkyl which is unsubstituted or substituted with  $\text{C}_3$ - $\text{C}_8$  cycloalkyl or  $\text{Ar}_1$ ;

10

$\text{X}_2$  is O or  $\text{NR}_5$ , wherein  $\text{R}_5$  is selected from the group consisting of hydrogen,  $\text{C}_1$ - $\text{C}_6$  straight or branched chain alkyl, and  $\text{C}_2$ - $\text{C}_6$  straight or branched chain alkenyl; and

15

$\text{R}_4$  is selected from the group consisting of phenyl, benzyl,  $\text{C}_1$ - $\text{C}_5$  straight or branched chain alkyl,  $\text{C}_2$ - $\text{C}_5$  straight or branched chain alkenyl,  $\text{C}_1$ - $\text{C}_5$  straight or branched chain alkyl substituted with phenyl, and  $\text{C}_2$ - $\text{C}_5$  straight or branched chain alkenyl substituted with phenyl.

20

6. The method of claim 5, wherein Z and  $\text{R}_1$  are lipophilic.

25

7. The method of claim 5, wherein the compound is selected from the group consisting of:

3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(3,4,5-trimethoxyphenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

5 3-(3,4,5-trimethoxyphenyl)-1-prop-2-(E)-enyl  
(2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(4,5-dichlorophenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

10 3-(4,5-dichlorophenyl)-1-prop-2-(E)-enyl (2S)-1-  
(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(4,5-methylenedioxyphenyl)-1-propyl (2S)-1-  
(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

15 3-(4,5-methylenedioxyphenyl)-1-prop-2-(E)-enyl  
(2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-cyclohexyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

20 3-cyclohexyl-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

(1R)-1,3-diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

25 (1R)-1,3-diphenyl-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

(1R)-1-cyclohexyl-3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;



(1R)-1-cyclohexyl-3-phenyl-1-prop-2-(E)-enyl  
 (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

(1R)-1-(4,5-dichlorophenyl)-3-phenyl-1-propyl  
 5 (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2S)-1-(1,2-dioxo-2-cyclohexyl)ethyl-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2S)-1-(1,2-dioxo-4-cyclohexyl)butyl-2-pyrrolidinecarboxylate;  
 10

3-phenyl-1-propyl (2S)-1-(1,2-dioxo-2-[2-furanyl])ethyl-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2S)-1-(1,2-dioxo-2-[2-thienyl])ethyl-2-pyrrolidinecarboxylate;

15 3-phenyl-1-propyl (2S)-1-(1,2-dioxo-2-[2-thiazolyl])ethyl-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2S)-1-(1,2-dioxo-2-phenyl)ethyl-2-pyrrolidinecarboxylate;

1,7-diphenyl-4-heptyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;  
 20

3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxo-4-hydroxybutyl)-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxamide;

25 1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylalanine ethyl ester;

1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-leucine ethyl ester;

1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylglycine ethyl ester;

1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylalanine phenyl ester;


5 1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylalanine benzyl ester;

1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-isoleucine ethyl ester; and

10 pharmaceutically acceptable salts, esters, and solvates thereof.

8. The method of claim 1, wherein the pyrrolidine derivative is a compound of formula II

15



II

20


or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

25 R<sub>1</sub> is C<sub>1</sub>-C<sub>8</sub>, straight or branched chain alkyl, C<sub>2</sub>-C<sub>8</sub>, straight or branched chain alkenyl, C<sub>3</sub>-C<sub>8</sub>, cycloalkyl, C<sub>5</sub>-C<sub>7</sub>, cycloalkenyl or Ar<sub>1</sub>, wherein said R<sub>1</sub> is unsubstituted or substituted with one or more substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>3</sub>-C<sub>8</sub>

cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, hydroxy, and Ar<sub>2</sub>;

Ar<sub>1</sub> and Ar<sub>2</sub> are independently selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl and phenyl, wherein said Ar<sub>1</sub> is unsubstituted or substituted with one or more substituent(s) independently selected from the group consisting of hydrogen, halo, hydroxy, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, and amino;

Z is C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, wherein said Z is substituted with one or more substituent(s) independently selected from the group consisting of Ar<sub>1</sub>, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl; or Z is fragment



wherein:

R<sub>3</sub> is C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl which is unsubstituted or substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl or Ar<sub>1</sub>;


$X_2$  is O or  $NR_5$ , wherein  $R_5$  is selected from the group consisting of hydrogen,  $C_1-C_6$  straight or branched chain alkyl, and  $C_2-C_6$  straight or branched chain alkenyl; and

5  $R_4$  is selected from the group consisting of phenyl, benzyl,  $C_1-C_5$  straight or branched chain alkyl,  $C_2-C_5$  straight or branched chain alkenyl,  $C_1-C_5$  straight or branched chain alkyl substituted with phenyl, and  $C_2-C_5$  straight or branched chain alkenyl substituted with phenyl.

9. The method of claim 8, wherein  $R_1$  is selected from the group consisting of  $C_1-C_6$  straight or branched chain alkyl, 2-cyclohexyl, 4-cyclohexyl, 2-furanyl, 2-thienyl, 2-thiazolyl, and 4-hydroxybutyl.

10. The method of claim 8, wherein Z and  $R_1$  are lipophilic.


20 11. The method of claim 1, wherein the pyrrolidine derivative is a compound of formula III



III

or a pharmaceutically acceptable salt, ester, or solvate or hydrate thereof, wherein:

Z' is fragment



wherein:

10 R<sub>3</sub> is C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl or unsubstituted Ar<sub>1</sub>, wherein said alkyl is unsubstituted or substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl or Ar<sub>1</sub>;


15 X<sub>2</sub> is O or NR<sub>5</sub>, wherein R<sub>5</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, and C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl;

20 R<sub>4</sub> is selected from the group consisting of phenyl, benzyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl substituted with phenyl, and C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl substituted with phenyl; and

Ar<sub>1</sub> is as defined in claim 8.

25 12. The method of claim 11, wherein Z' is lipophilic.

13. The method of claim 1, wherein the pyrrolidine derivative is a compound of formula IV



IV

5

or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

10  $R_1$  is  $C_1$ - $C_6$  straight or branched chain alkyl,  $C_2$ - $C_6$  straight or branched chain alkenyl,  $C_3$ - $C_6$  cycloalkyl or  $Ar_1$ , wherein said alkyl or alkenyl is unsubstituted or substituted with  $C_3$ - $C_6$  cycloalkyl or  $Ar_2$ ;

$Ar_1$  and  $Ar_2$  are independently selected from the group consisting of 2-furyl, 2-thienyl, and phenyl;

15  $X$  is selected from the group consisting of oxygen and sulfur;

$Y$  is oxygen;

20  $Z$  is  $C_1$ - $C_6$  straight or branched chain alkyl, or  $C_2$ - $C_6$  straight or branched chain alkenyl, wherein said  $Z$  is substituted with one or more substituent(s) independently selected from the group consisting of 2-furyl, 2-thienyl,  $C_3$ - $C_6$  cycloalkyl, pyridyl, and phenyl, each having one or more substituent(s) independently selected from the group consisting of  
25 hydrogen and  $C_1$ - $C_4$  alkoxy.

14. The method of claim 13, wherein  $Z$  and  $R_1$  are lipophilic.

15. The method of claim 13, wherein the compound is selected from the group consisting of:

3-(2,5-dimethoxyphenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

5 3-(2,5-dimethoxyphenyl)-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

2-(3,4,5-trimethoxyphenyl)-1-ethyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

10 3-(3-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(2-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(4-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

15 3-phenyl-1-propyl (2S)-1-(2-tert-butyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2S)-1-(2-cyclohexylethyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

20 3-(3-pyridyl)-1-propyl (2S)-1-(2-cyclohexylethyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

3-(3-pyridyl)-1-propyl (2S)-1-(2-tert-butyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

3,3-diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

25 3-(3-pyridyl)-1-propyl (2S)-1-(2-cyclohexyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

3-(3-pyridyl)-1-propyl (2S)-N-([2-thienyl]

glyoxyl)pyrrolidinecarboxylate;

3,3-diphenyl-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-dioxobutyl)-2-pyrrolidinecarboxylate;

3,3-diphenyl-1-propyl (2*S*)-1-cyclohexylglyoxyl-  
5 2-pyrrolidinecarboxylate;

3,3-diphenyl-1-propyl (2*S*)-1-(2-thienyl)glyoxyl-  
2-pyrrolidinecarboxylate; and

pharmaceutically acceptable salts, esters, and  
solvates thereof.

10

16. The method of claim 15, wherein the compound  
is selected from the group consisting of:

3-(3-pyridyl)-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-  
dioxopentyl)-2-pyrrolidinecarboxylate;

15 3-(2-pyridyl)-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-  
dioxopentyl)-2-pyrrolidinecarboxylate;

3-(3-pyridyl)-1-propyl (2*S*)-1-(2-cyclohexyl-1,2-  
dioxoethyl)-2-pyrrolidinecarboxylate; and


20 pharmaceutically acceptable salts, esters, and  
solvates thereof.

17. The method of claim 16, wherein the compound  
is 3-(3-pyridyl)-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-  
25 dioxopentyl)-2-pyrrolidinecarboxylate, or a  
pharmaceutically acceptable salt, ester, or solvate or  
hydrate thereof.



18. The method of claim 1, wherein the pyrrolidine derivative is an N-glyoxyl prolyl ester.

19. The method of claim 1, wherein the  
5 pyrrolidine derivative is a compound of formula V



V

or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

15 V is C, N, or S;

A and B, taken together with V and the carbon atom to which they are respectively attached, form a 5-7 membered saturated or unsaturated heterocyclic ring containing, in addition to V, one or more  
20 heteroatom(s) selected from the group consisting of O, S, SO, SO<sub>2</sub>, N, NH, and NR;

R is either C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>9</sub> straight or branched chain alkenyl, C<sub>3</sub>-C<sub>9</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, or Ar<sub>1</sub>, wherein R is  
25 either unsubstituted or substituted with one or more substituent(s) independently selected from the group consisting of halo, haloalkyl, carbonyl, carboxy, hydroxy, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or

alkenyl,  $C_1-C_4$  alkoxy,  $C_2-C_4$  alkenyloxy, phenoxy, benzyloxy, thioalkyl, alkylthio, sulfhydryl, amino, alkylamino, aminoalkyl, aminocarboxyl, and  $Ar_2$ ;

5  $R_1$  is  $C_1-C_9$  straight or branched chain alkyl,  $C_2-C_9$  straight or branched chain alkenyl,  $C_3-C_8$  cycloalkyl,  $C_5-C_7$  cycloalkenyl or  $Ar_1$ , wherein said  $R_1$  is unsubstituted or substituted with one or more substituents independently selected from the group  
10 consisting of  $C_1-C_6$  alkyl,  $C_2-C_6$  alkenyl,  $C_3-C_8$  cycloalkyl,  $C_5-C_7$  cycloalkenyl, hydroxy, and  $Ar_2$ ;

$Ar_1$  and  $Ar_2$  are independently an alicyclic or aromatic, mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either  
15 unsubstituted or substituted with one or more substituent(s); wherein the individual ring size is 5-8 members; wherein said heterocyclic ring contains 1-6 heteroatom(s) independently selected from the group consisting of O, N, and S;


20 X is O, S,  $CH_2$  or  $H_2$ ;

Y is O or  $NR_2$ , wherein  $R_2$  is hydrogen or  $C_1-C_6$  alkyl; and

Z is  $C_1-C_6$  straight or branched chain alkyl, or  $C_2-C_6$  straight or branched chain alkenyl, wherein said  
25 Z is substituted with one or more substituent(s) independently selected from the group consisting of  $Ar_1$ ,  $C_3-C_8$  cycloalkyl, and  $C_1-C_6$  straight or branched chain alkyl or  $C_2-C_6$  straight or branched chain alkenyl

substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl; or Z is fragment

5



wherein:

10

R<sub>3</sub> is C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl which is unsubstituted or substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl or Ar<sub>1</sub>;

X<sub>2</sub> is O or NR<sub>5</sub>, wherein R<sub>5</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, and C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl; and

15

R<sub>4</sub> is selected from the group consisting of phenyl, benzyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl substituted with phenyl, and C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl substituted with phenyl.

20

20. A pharmaceutical composition which comprises:

25


- (i) an effective amount of a pyrrolidine derivative for treating alopecia or promoting hair growth in an animal; and
- (ii) a pharmaceutically acceptable carrier.

21. The pharmaceutical composition of claim 20, wherein the pyrrolidine derivative is non-immunosuppressive.

5 22. The pharmaceutical composition of claim 20, wherein the pyrrolidine derivative has an affinity for an FKBP-type immunophilin.

23. The pharmaceutical composition of claim 22, 10 wherein the FKBP-type immunophilin is FKBP-12.

24. The pharmaceutical composition of claim 20, wherein the pyrrolidine derivative is a compound of formula I



or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

25  $R_1$  is  $C_1$ - $C_8$  straight or branched chain alkyl,  $C_2$ - $C_8$  straight or branched chain alkenyl,  $C_3$ - $C_8$  cycloalkyl,  $C_5$ - $C_7$  cycloalkenyl or  $Ar_1$ , wherein said  $R_1$  is unsubstituted or substituted with one or more substituents independently selected from the group

consisting of  $C_1-C_6$  alkyl,  $C_2-C_6$  alkenyl,  $C_3-C_8$  cycloalkyl,  $C_5-C_7$  cycloalkenyl, hydroxy, and  $Ar_2$ ;


$Ar_1$  and  $Ar_2$  are independently selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl and phenyl, wherein said  $Ar_1$  is unsubstituted or substituted with one or more substituent(s) independently selected from the group consisting of hydrogen, halo, hydroxy, nitro, trifluoromethyl,  $C_1-C_6$  straight or branched chain alkyl,  $C_2-C_6$  straight or branched chain alkenyl,  $C_1-C_4$  alkoxy,  $C_2-C_4$  alkenyloxy, phenoxy, benzyloxy, and amino;

X is O, S,  $CH_2$  or  $H_2$ ;

Y is O or  $NR_2$ , wherein  $R_2$  is hydrogen or  $C_1-C_6$  alkyl; and

Z is  $C_1-C_6$  straight or branched chain alkyl, or  $C_2-C_6$  straight or branched chain alkenyl, wherein said Z is substituted with one or more substituent(s) independently selected from the group consisting of  $Ar_1$ ,  $C_3-C_8$  cycloalkyl, and  $C_1-C_6$  straight or branched chain alkyl or  $C_2-C_6$  straight or branched chain alkenyl substituted with  $C_3-C_8$  cycloalkyl; or Z is fragment

25



wherein:

$R_3$  is  $C_1$ - $C_9$  straight or branched chain alkyl which is unsubstituted or substituted with  $C_3$ - $C_8$  cycloalkyl or  $Ar_1$ ;

5  $X_2$  is O or  $NR_5$ , wherein  $R_5$  is selected from the group consisting of hydrogen,  $C_1$ - $C_6$  straight or branched chain alkyl, and  $C_2$ - $C_6$  straight or branched chain alkenyl; and

10  $R_4$  is selected from the group consisting of phenyl, benzyl,  $C_1$ - $C_5$  straight or branched chain alkyl,  $C_2$ - $C_5$  straight or branched chain alkenyl,  $C_1$ - $C_5$  straight or branched chain alkyl substituted with phenyl, and  $C_2$ - $C_5$  straight or branched chain alkenyl substituted with phenyl.

15

25. The pharmaceutical composition of claim 24, wherein Z and  $R_1$  are lipophilic.

20 26. The pharmaceutical composition of claim 24, wherein the compound is selected from the group consisting of:

3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

25 3-phenyl-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(3,4,5-trimethoxyphenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(3,4,5-trimethoxyphenyl)-1-prop-2-(E)-enyl

(2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(4,5-dichlorophenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

5 3-(4,5-dichlorophenyl)-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(4,5-methylenedioxyphenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

10 3-(4,5-methylenedioxyphenyl)-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-cyclohexyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

15 3-cyclohexyl-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

(1R)-1,3-diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

20 (1R)-1,3-diphenyl-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

(1R)-1-cyclohexyl-3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

(1R)-1-cyclohexyl-3-phenyl-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

25 (1R)-1-(4,5-dichlorophenyl)-3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-

pyrrolidinecarboxylate;



3-phenyl-1-propyl (2*S*)-1-(1,2-dioxo-2-cyclohexyl)ethyl-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2*S*)-1-(1,2-dioxo-4-cyclohexyl)butyl-2-pyrrolidinecarboxylate;

5 3-phenyl-1-propyl (2*S*)-1-(1,2-dioxo-2-[2-furanyl])ethyl-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2*S*)-1-(1,2-dioxo-2-[2-thienyl])ethyl-2-pyrrolidinecarboxylate;

10 3-phenyl-1-propyl (2*S*)-1-(1,2-dioxo-2-[2-thiazolyl])ethyl-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2*S*)-1-(1,2-dioxo-2-phenyl)ethyl-2-pyrrolidinecarboxylate;

1,7-diphenyl-4-heptyl (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

15 3-phenyl-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-dioxo-4-hydroxybutyl)-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2*S*)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxamide;


20 1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylalanine ethyl ester;

1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-leucine ethyl ester;

1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylglycine ethyl ester;

25 1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylalanine phenyl ester;

1-[1-(3,3-dimethyl-1,2-dioxopentyl)-L-proline]-L-phenylalanine benzyl ester;



II

15


or a pharmaceutically acceptable salt thereof,  
wherein:

20  $R_1$  is  $C_1$ - $C_6$  straight or branched chain alkyl,  $C_2$ - $C_6$  straight or branched chain alkenyl,  $C_3$ - $C_8$  cycloalkyl,  $C_5$ - $C_7$  cycloalkenyl or  $Ar_1$ , wherein said  $R_1$  is unsubstituted or substituted with one or more substituents independently selected from the group consisting of  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_3$ - $C_8$  cycloalkyl,  $C_5$ - $C_7$  cycloalkenyl, hydroxy, and  $Ar_2$ ;

25  $Ar_1$  and  $Ar_2$  are independently selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl and phenyl, wherein said

Ar<sub>1</sub> is unsubstituted or substituted with one or more substituent(s) independently selected from the group consisting of hydrogen, halo, hydroxy, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, and amino;

Z is C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, wherein said Z is substituted with one or more substituent(s) independently selected from the group consisting of Ar<sub>1</sub>, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl; or Z is fragment



wherein:

R<sub>3</sub> is C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl which is unsubstituted or substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl or Ar<sub>1</sub>;

X<sub>2</sub> is O or NR<sub>5</sub>, wherein R<sub>5</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, and C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl; and


R<sub>4</sub> is selected from the group consisting of

phenyl, benzyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl substituted with phenyl, and C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl substituted with phenyl.

28. The pharmaceutical composition of claim 27, wherein R<sub>1</sub> is selected from the group consisting of C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl, 2-cyclohexyl, 4-cyclohexyl, 2-furanyl, 2-thienyl, 2-thiazolyl, and 4-hydroxybutyl.

29. The pharmaceutical composition of claim 27, wherein Z and R<sub>1</sub> are lipophilic.


30. The pharmaceutical composition of claim 20, wherein the pyrrolidine derivative is a compound of formula III



III

or a pharmaceutically acceptable salt, ester, or solvate or hydrate thereof, wherein:

Z is fragment



5

wherein:

$\text{R}_3$  is  $\text{C}_1\text{-C}_6$  straight or branched chain alkyl or unsubstituted  $\text{Ar}_1$ , wherein said alkyl is unsubstituted or substituted with  $\text{C}_3\text{-C}_8$  cycloalkyl or  $\text{Ar}_1$ ;

10

$\text{X}_2$  is O or  $\text{NR}_5$ , wherein  $\text{R}_5$  is selected from the group consisting of hydrogen,  $\text{C}_1\text{-C}_6$  straight or branched chain alkyl, and  $\text{C}_2\text{-C}_6$  straight or branched chain alkenyl;

15

$\text{R}_4$  is selected from the group consisting of phenyl, benzyl,  $\text{C}_1\text{-C}_6$  straight or branched chain alkyl,  $\text{C}_2\text{-C}_6$  straight or branched chain alkenyl,  $\text{C}_1\text{-C}_6$  straight or branched chain alkyl substituted with phenyl, and  $\text{C}_2\text{-C}_6$  straight or branched chain alkenyl substituted with phenyl; and


20

$\text{Ar}_1$  is as defined in claim 26.

31. The pharmaceutical composition of claim 30, wherein  $\text{Z}'$  is lipophilic.

25

32. The pharmaceutical composition of claim 20, wherein the pyrrolidine derivative is a compound of formula IV



IV

5

or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

10  $R_1$  is  $C_1$ - $C_6$  straight or branched chain alkyl,  $C_2$ - $C_6$  straight or branched chain alkenyl,  $C_3$ - $C_6$  cycloalkyl or  $Ar_1$ , wherein said alkyl or alkenyl is unsubstituted or substituted with  $C_3$ - $C_6$  cycloalkyl or  $Ar_2$ ;

$Ar_1$  and  $Ar_2$  are independently selected from the group consisting of 2-furyl, 2-thienyl, and phenyl;

15  $X$  is selected from the group consisting of oxygen and sulfur;

$Y$  is oxygen;

20  $Z$  is  $C_1$ - $C_6$  straight or branched chain alkyl, or  $C_2$ - $C_6$  straight or branched chain alkenyl, wherein said  $Z$  is substituted with one or more substituent(s) independently selected from the group consisting of 2-furyl, 2-thienyl,  $C_3$ - $C_6$  cycloalkyl, pyridyl, and phenyl, each having one or more substituent(s) independently selected from the group consisting of

25 hydrogen and  $C_1$ - $C_4$  alkoxy.

33. The pharmaceutical composition of claim 32, wherein  $Z$  and  $R_1$  are lipophilic.

34. The pharmaceutical composition of claim 32, wherein the compound is selected from the group consisting of:

3-(2,5-dimethoxyphenyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(2,5-dimethoxyphenyl)-1-prop-2-(E)-enyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

2-(3,4,5-trimethoxyphenyl)-1-ethyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(3-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(2-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(4-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2S)-1-(2-tert-butyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

3-phenyl-1-propyl (2S)-1-(2-cyclohexylethyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

3-(3-pyridyl)-1-propyl (2S)-1-(2-cyclohexylethyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

3-(3-pyridyl)-1-propyl (2S)-1-(2-tert-butyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

3,3-diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(3-pyridyl)-1-propyl (2S)-1-(2-cyclohexyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate;

glyoxyl)pyrrolidinecarboxylate;

3,3-diphenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxobutyl)-2-pyrrolidinecarboxylate;

5        3,3-diphenyl-1-propyl (2S)-1-cyclohexylglyoxyl-2-pyrrolidinecarboxylate;

3,3-diphenyl-1-propyl (2S)-1-(2-thienyl)glyoxyl-2-pyrrolidinecarboxylate; and

10        pharmaceutically acceptable salts, esters, and solvates thereof.

35. The pharmaceutical composition of claim 34, wherein the compound is selected from the group consisting of:

15        3-(3-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

3-(2-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate;

20        3-(3-pyridyl)-1-propyl (2S)-1-(2-cyclohexyl-1,2-dioxoethyl)-2-pyrrolidinecarboxylate; and

pharmaceutically acceptable salts, esters, and solvates thereof.


36. The pharmaceutical composition of claim 35, wherein the compound is 3-(3-pyridyl)-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate, or a pharmaceutically acceptable salt, ester, or solvate or hydrate thereof.

25



37. The pharmaceutical composition of claim 20, wherein the pyrrolidine derivative is an N-glyoxyl prolyl ester.

38. The pharmaceutical composition of claim 20, wherein the pyrrolidine derivative is a compound of formula V



or a pharmaceutically acceptable salt, ester, or solvate thereof, wherein:

V is C, N, or S;

A and B, taken together with V and the carbon atom to which they are respectively attached, form a 5-7 membered saturated or unsaturated heterocyclic ring containing, in addition to V, one or more heteroatom(s) selected from the group consisting of O, S, SO, SO<sub>2</sub>, N, NH, and NR;

R is either C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>9</sub> straight or branched chain alkenyl, C<sub>3</sub>-C<sub>9</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkenyl, or Ar<sub>1</sub>, wherein R is either unsubstituted or substituted with one or more substituent(s) independently selected from the group

consisting of halo, haloalkyl, carbonyl, carboxy, hydroxy, nitro, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>2</sub>-C<sub>4</sub> alkenyloxy, phenoxy, benzyloxy, thioalkyl, alkylthio, sulfhydryl, amino, alkylamino, aminoalkyl, aminocarboxyl, and Ar<sub>2</sub>;

R<sub>1</sub> is C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>9</sub> straight or branched chain alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl or Ar<sub>1</sub>, wherein said R<sub>1</sub> is unsubstituted or substituted with one or more substituents independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>5</sub>-C<sub>7</sub> cycloalkenyl, hydroxy, and Ar<sub>2</sub>;

Ar<sub>1</sub> and Ar<sub>2</sub> are independently an alicyclic or aromatic, mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either unsubstituted or substituted with one or more substituent(s); wherein the individual ring size is 5-8 members; wherein said heterocyclic ring contains 1-6 heteroatom(s) independently selected from the group consisting of O, N, and S;


X is O, S, CH<sub>2</sub> or H<sub>2</sub>;

Y is O or NR<sub>2</sub>, wherein R<sub>2</sub> is hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl; and

Z is C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl, wherein said Z is substituted with one or more substituent(s) independently selected from the group consisting of

Ar<sub>1</sub>, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl or C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl; or Z is fragment

5



wherein:

10

R<sub>3</sub> is C<sub>1</sub>-C<sub>9</sub> straight or branched chain alkyl which is unsubstituted or substituted with C<sub>3</sub>-C<sub>8</sub> cycloalkyl or Ar<sub>1</sub>;

15

X<sub>2</sub> is O or NR<sub>5</sub>, wherein R<sub>5</sub> is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> straight or branched chain alkyl, and C<sub>2</sub>-C<sub>6</sub> straight or branched chain alkenyl; and

20

R<sub>4</sub> is selected from the group consisting of phenyl, benzyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl, C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl, C<sub>1</sub>-C<sub>5</sub> straight or branched chain alkyl substituted with phenyl, and C<sub>2</sub>-C<sub>5</sub> straight or branched chain alkenyl substituted with phenyl.

1 / 5

FIG. 1



FIG.2




FIG.3



FIG.4



**FIG. 5**  
Promotion of Hair Growth by Neuroimmunophilin Ligands





## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K7/48 A61K31/40 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages                                      | Relevant to claim No. |
|------------|---|-----------------------|
| X          | EP 0 423 714 A (FUJISAWA PHARM. CO., LTD)<br>24 April 1991<br>cited in the application<br>see the whole document<br>--- | 1-38                  |
| X          | WO 93 18736 A (A PARDO ET AL.)<br>30 September 1993<br>see the whole document<br>---                                    | 1-4,<br>20-23         |
| X          | EP 0 519 819 A (L'OREAL) 23 December 1992<br><br>see claims 1,16-18; example 5<br>---                                   | 1-4,<br>20-23         |
| P, X       | US 5 714 510 A (P. PROCTOR)<br>3 February 1998<br>see the whole document<br>---   | 1-4,<br>20-23         |
| -/--       |   |                       |

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

14 October 1998

Date of mailing of the international search report

21/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Glikman, J-F

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category | Citation of document, with indication, where appropriate, of the relevant passages       | Relevant to claim No. |
|----------|--|-----------------------|
| P,X      | WO 98 13343 A (GUILFORD PHARM., INC.)<br>2 April 1998<br>see the whole document<br>----- | 20-38                 |

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1-38 (PARTIAL)  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
In view of the large number of compounds which are defined by the wording of the claims, the search has been performed on the general idea and compounds mentioned in the examples of the description.
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

| Patent document<br>cited in search report |   | Publication<br>date | Patent family<br>member(s)   | Publication<br>date  |
|---|---|---------------------|--|--|
| EP 423714                                 | A | 24-04-1991          | AT 107499 T<br>CA 2027608 A<br>DE 69010139 D<br>DE 69010139 T<br>DK 423714 T<br>JP 3204807 A<br>US 5215995 A   | 15-07-1994<br>17-04-1991<br>28-07-1994<br>13-10-1994<br>25-07-1994<br>06-09-1991<br>01-06-1993   |
| WO 9318736                                | A | 30-09-1993          | NONE   |  |
| EP 519819                                 | A | 23-12-1992          | FR 2677884 A<br>AT 125148 T<br>CA 2071802 A<br>DE 69203545 D<br>DE 69203545 T<br>ES 2075648 T<br>JP 5186315 A<br>US 5772990 A  | 24-12-1992<br>15-08-1995<br>21-12-1992<br>24-08-1995<br>11-01-1996<br>01-10-1995<br>27-07-1993<br>30-06-1998   |
| US 5714510                                | A | 03-02-1998          | US 5470876 A<br>US 5472687 A<br>US 5352442 A<br>US 5714482 A<br>US 5716947 A<br>US 5723502 A<br>US 5728714 A<br>AT 110954 T<br>DE 68917946 D<br>DE 68917946 T<br>EP 0327263 A<br>AU 1362488 A<br>WO 8805653 A<br>AU 6131686 A<br>DE 3688427 D<br>EP 0232311 A<br>WO 8700427 A<br>JP 63500865 T | 28-11-1995<br>05-12-1995<br>04-10-1994<br>03-02-1998<br>10-02-1998<br>03-03-1998<br>17-03-1998<br>15-09-1994<br>13-10-1994<br>05-01-1995<br>09-08-1989<br>24-08-1988<br>11-08-1988<br>10-02-1987<br>17-06-1993<br>19-08-1987<br>29-01-1987<br>31-03-1988 |
| WO 9813343                                | A | 02-04-1998          | US 5786378 A<br>AU 4259097 A   | 28-07-1998<br>17-04-1998   |